



UNITED NATIONS

E/NL. 1957/117

3 December 1957

Original : ENGLISH

LAWS AND REGULATIONS

PROMULGATED TO GIVE EFFECT TO THE PROVISIONS OF THE CONVENTION OF 13 JULY 1931 FOR LIMITING THE MANUFACTURE AND REGULATING THE DISTRIBUTION OF NARCOTIC DRUGS, AS AMENDED BY THE PROTOCOL OF 11 DECEMBER 1946

FEDERATION OF RHODESIA AND NYASALAND

Communicated by the Government of the United Kingdom of Great Britain and Northern Ireland

NOTE BY THE SECRETARY-GENERAL-- In accordance with Article 21 of the Convention of 13 July 1931 for Limiting the Manufacture and Regulating the Distribution of Narcotic Drugs, as amended by the Protocol of 11 December 1946, the Secretary-General has the honour to communicate the following legislative text.

Supplement to the Federal Government Gazette,
16 March 1956

ACT 28/55

FEDERAL GOVERNMENT NOTICE No. 58 OF 1956.

DECLARATION IN TERMS OF SECTION 10 (4) OF THE DANGEROUS DRUGS ACT, 1955¹⁾

It is hereby notified that the Minister of Health has, in terms of subsection (4) of section 10 of the Dangerous Drugs Act, 1955, declared that the provisions of Part III of that Act shall not apply to the following preparation after the 31st day of March, 1956.

A. - MORPHINE PREPARATIONS:

	Substance	Formula
1. Cereoli iodoformi et morphinae	Iodoform Morphine Hydrochloride Oil of theobroma, sufficient to fill a 1 - grammé mould.	0.320 grammé 0.016 grammé sufficient to fill a 1 - grammé mould.
2. Emplastrum opii	Elemi Terebinthina Cera flava Olibanum pulvis Benzoes pulvis Opii pulvis Balsamum peruvianum	20 grammes 30 grammes 15 grammes 18 grammes 10 grammes 5 grammes 2 grammes
3. Emplastrum opii	Extract of opium Refined elemi Diachylon plaster with gum . . .	25 grammes 25 grammes 50 grammes

1) Note by the Secretariat: E/NL.1956/44

Substance	Formula
4. Emplastrum opii	Elemi 8 grammes Terebinthinae communis 15 grammes Ceroe flavae 5 grammes Olibani pulveratae 8 grammes Benzoes pulveratae 4 grammes Opii pulverati 2 grammes Balsami peruviani 1 gramme
5. Emplastrum opii	Opium, in very fine powder 10 grammes Resin plaster 90 grammes
6. Emplastrum opii (see formula under 5) mixed with other plasters contained in the British Pharmacopoeia or British Pharmaceutical Codex.	
7. Linimentum opii	Tincture of opium 500 millilitres Liniment of soap 500 millilitres
8. Linimentum opii (see formula under 7) mixed with any other liniment of the British Pharmacopoeia or British Pharmaceutical Codex.	
9. Linimentum opii ammoniatum	Ammoniated liniment of camphor 30 Tincture of opium 30 Liniment of Belladonna 5 Strong solution of ammonia 5 Liniment of soap to 100
10. Linimentum opii ammoniatum (see formula under 9) mixed with any other British Pharmacopoeia or British Pharmaceutical Codex liniment.	
11. Caustic "Nerve Pastes"	Preparations containing, in addition to morphine salts, or morphine and cocaine salts, at least 25 per centum of arsenious acid, and made up with the requisite proportion of creosote or phenol to produce the consistency of a paste.
12. Diarrhoea pills	Camphor 0.0648 gramme Lead acetate 0.013 gramme Bismuth subnitrate 0.162 gramme Tannic acid 0.0648 gramme Opium powder 0.026 gramme
13. Pilulae digitalis et Opii compositae	Digitalis leaves, in powder 0.31 gramme Opium, in powder 0.19 gramme Ipecacuanha root, in powder 0.13 gramme Quinine sulphate 0.78 gramme Syrup of glucose, a sufficient quantity to make 12 pills.
14. Pilulae hydrargyri cum Opio	Mercury pill 3.89 grammes Opium in powder 0.19 gramme To make 12 pills.

Substance	Formula
15. Pilulae hydrargyri cum Creta et Opii	Mercury with chalk 0.78 grammie Compound powder of ipecacuanha* . 0.78 grammie Milk sugar, a sufficient quantity Syrup of glucose, a sufficient quantity To make 12 pills.
16. Pilulae ipecacuanhae cum Scilla	Compound powder of ipecacuanha* . 30 grammes Squill, in powder 10 grammes Ammoniacum, in powder 10 grammes Syrup of glucose, a sufficient quantity.
17. Pilulae hydrargyri bichlorati cum Opii extracto.	Bichloride of mercury triturated . . 10 centigrammes Extract of opium 20 centigrammes Extract of couchgrass 20 centigrammes Liquorice root in powder, q.s. for 10 pills.
18. Pilulae hydrargyri iodati cum Opii pulvere . . .	Hydrargyrum iodatum freshly prepared 50 centigrammes Opium powder 20 centigrammes Powdered Liquorice 30 centigrammes White honey, q.s. for 10 pills.
19. Pilula plumbi cum Opio.	Lead acetate, in powder 80 grammes Opium, in powder 12 grammes Syrup of glucose 8 grammes (or a sufficient quantity).
20. Pilulae terebinthinae compositae	Opium 0.5 grammie Chinina sulfas 2 grammes Styrax liquidus 2 grammes Terebinthina laricina 8 grammes Magnesii subcarbonas, a sufficient quantity to make 100 pills.
21. Pulvis Ipecacuanhae et Opii (Dover's Powder) .	Powdered Ipecacuanha 100 grammes Powdered Opium 100 grammes Potassium sulphate in powder 800 grammes
22. Mixtures of Dover's Powder (see formula under 21) with mercury and chalk, aspirin, phenacetin, quinine and its salts, and sodium bicarbonate.	
23. Pulvis kino compositus	Kino, in powder 75 grammes Opium, in powder 5 grammes Cinnamon bark, in powder 20 grammes
24. Suppositoria plumbi composita Syn.: Suppositoria plumbi cum opio	Lead acetate, in powder 2.4 grammes Opium, in powder 0.8 grammie Oil of theobroma, a sufficient quantity for 12 sup- positories, each weighing about 1 grammie.
25. Coryza Tablets, No. 2	Powdered opium 0.0043 grammie Quinine sulph. 0.022 grammie Ammon. chlor. 0.022 grammie Camphor 0.022 grammie Ext. belladonna leaves. 0.0043 grammie Ext. aconite root. 0.0043 grammie

* The formula of this powder is given under 21.

Substance	Formula
26. Diarrhoea Tablets, No. 2	
Powdered opium	0.016 grammæ
Camphor	0.016 grammæ
Powdered ipecacuanha	0.008 grammæ
Lead acetate	0.011 grammæ
27. Dysentery Tablets	
Powdered opium	0.013 grammæ
Powdered ipecacuanha	0.0648 grammæ
Powdered calomel	0.0324 grammæ
Lead acetate	0.0324 grammæ
Bismuth betanaphthol	0.1944 grammæ
28. Tabella hydrargyri cum Opio	
Mercurous chloride powder	0.065 grammæ
Antimony oxide powder	0.065 grammæ
Ipecacuanha-root powder	0.065 grammæ
Powdered opium	0.065 grammæ
Milk sugar	0.065 grammæ
Gelatine solution, a sufficient quantity to make one tablet.	
29. Tabella plumbi cum Opio	
Sugar of lead	0.195 grammæ
Powdered opium	0.065 grammæ
Gelatine solution, a sufficient quantity to make 1 tablet.	
30. Tablettæ plumbi cum Opio	
Lead acetate, in fine powder	19.44 grammes
Opium, in powder	3.24 grammes
Refined sugar, in powder	6.48 grammes
Ethereal solution of theobroma	3.60 mils.
Alcohol	0.90 mil.
31. Unguentum gallæ compositum	
Galls in very fine powder	20
Extract of opium	4
Distilled water	16
Wool fat	10
Soft paraffin, yellow	50
32. Unguentum gallæ compositum (see formula under 31) mixed with other ointment and plasters contained in the British Pharmacopœia or British Pharmaceutical Codex.	
33. Unguentum gallæ cum Opio	
Gall ointment	92.5 grammes
Opium in powder	7.5 grammes
34. Unguentum gallæ cum opio (see formula under 33) mixed with other ointments and plasters contained in the British Pharmacopœia or British Pharmaceutical Codex.	
35. Yatren-105 (Iodoxyquinoline-sulphonic acid) with 5 per centum opium admixture.	

B. - COCAINE PREPARATIONS:

- | | | |
|-------------------------------------|--------------------------------------|--------------|
| 1. Bernatzik's Injections | (a) Hydrargyrum bicanatum | 0.03 grammes |
| | Cocainum | 0.02 grammes |
| | (b) Hydrargyrum succinatum | 0.03 grammes |
| | Cocainum | 0.01 grammæ |

Substance	Formula
2. Stila's Injections	(a) Hydrargyrum succinatum . . 0.03 grammes Cocainum muriaticum. . . . 0.01 grammes (b) Hydrargyrum succinatum . . 0.05 grammes Cocainum muriaticum . . . 0.03 grammes
3. Natrium biboracicum compositum cum Cocaino	In tablets, compressed tablets, lozenges, pastilles and the like, difficult to break up, and containing not more than 0.2 per centum of cocaine salts in conjunction with not less than 20 per centum borax and not less than 20 per centum antipyrine, or some similar analgesic, and not more than 40 per centum of flavouring matter. Maximum weight of each tablet, etc., 1 gramme.
4. Caustic "Nerve Pastes"	Preparations containing, in addition to cocaine salts or cocaine and morphine salts, at least 25 per centum of arsenious acid, and made up with the requisite proportion of creosote or phenol to produce the consistency of a paste.
5. Cocaine and Atropine Tablets, with a content of not more than 0.0003 grammes of cocaine salts and not less than 0.0003 grammes of atropine salts to each tablet.	Atropinum sulphuricum 0.0003 grammes Cocainum hydrochloricum . . . 0.0003 grammes Mannite 0.003 grammes Weight of one tablet 0.0036 grammes Cocaine content, 8.3 per centum.

C. - DICODIDE PREPARATIONS:

1. Cardiazol-Dicodide Solutions	Solutions containing not less than 10 per centum of cardiazol and not more than 0.5 per centum of dicodide salts.
---	---

D. - EUCODAL PREPARATIONS:

1. Anti-Opium Tablets	Eucodal 1 grammes Pulvis gentianae 35 grammes Pulvis ipecacuanhae 20 grammes Quinine sulphate 20 grammes Caffeine 5 grammes Sugar of milk 25 grammes Mix up and make up 5-grain tablets.
2. Tablets B.B. Compound	Berberis vulgaris powder 0.0324 grammes Nux vomica 0.013 grammes Eucodal 0.0032 grammes Ipecacuanha 0.0648 grammes Rhubarb 0.013 grammes Pulvis cinnamoni compositus . 0.0324 grammes Aromatic chalk 0.0032 grammes