

**General Assembly**

Distr.: General
26 March 2013
English
Original: French

Sixty-seventh session

Agenda item 75 (a)

Oceans and the law of the sea**Letter dated 15 March 2013 from the Permanent Representative of Monaco to the United Nations addressed to the Secretary-General**

I have the honour to draw attention to the conclusions of the workshop held in Monaco from 11 to 13 November 2012 on the topic “Bridging the gap between ocean acidification impacts and economic valuation” organized by the Monaco Scientific Centre and the Marine Environment Laboratories of the International Atomic Energy Agency (see annex).

I would be grateful if this letter and its annex could be circulated as a document of the sixty-seventh session of the General Assembly under agenda item 75 (a): “Oceans and the law of the sea”.

(Signed) Isabelle Picco

* Reissued for technical reasons on 23 April 2013.



Annex to the letter dated 15 March 2013 from the Permanent Representative of Monaco to the United Nations addressed to the Secretary-General

[Original: English and French]

Economics of ocean acidification

Impacts on fisheries and aquaculture

Ocean acidification has only recently been recognized as a threat for the environment with potentially severe social and economic consequences. The following statements are the major conclusions and recommendations from the second international workshop on “Bridging the gap between ocean acidification impacts and economic valuation”. The objective of the workshop was to assess ocean acidification impacts on fisheries and aquaculture resources in different regions of the world. The regions were artificial aggregations of the fishing areas defined by the Food and Agriculture Organization of the United Nations (FAO). Each region spanned multiple political, economic and ecological units.

General conclusions

- Ocean acidification is a global issue directly caused by increased anthropogenic carbon dioxide emissions to the atmosphere. It is happening now and carbon dioxide absorbed by the oceans will continue to rise long after those emissions are reduced.
- Some ocean areas, such as upwelling waters (deep water drawn upwards as wind pushes surface water offshore), polar and subpolar regions and some coastal and estuarine waters, are natural “hot spots” of special concern for ocean acidification.
- Coastal waters are also subject to warming, low oxygen, high nutrients and pollution in addition to ocean acidification, exposing aquatic species and communities to multiple stressors.
- World capture fisheries and aquaculture generated about \$218 billion in 2010. This provides an estimated 4.3 billion people with at least 15 per cent of their animal protein. Over the last 30 years, world food production by aquaculture has expanded twelvefold, to account for nearly half of human seafood consumption. Dependence on marine protein is expected to continue to rise with increased human population (FAO statistics).
- Major fisheries and aquaculture often occur in areas sensitive to ocean acidification. This puts economies and livelihoods at risk, and requires consideration and action by policymakers.
- Research on valuable seafood species is limited. Studies indicate that some shellfish used in aquaculture may be vulnerable to ocean acidification. Much uncertainty remains concerning finfish.

Recommendations

- Mitigate the effects of ocean acidification by reducing emissions of atmospheric carbon dioxide.
- Establish ad hoc coastal monitoring networks for standardized measurement of ocean acidification.
- Support research on high-value finfish, shrimp and other shellfish in high carbon dioxide conditions to enable socioeconomic assessment of impacts on food security.
- Implement best practices and adaptive management of fisheries and aquaculture to increase ecological resilience of marine ecosystems.
- Increase the adaptive capacity of fishing communities through education about ocean acidification and by training and support to diversify livelihoods where needed.
- Improve multi-stakeholder exchange of information and communication among parties (coastal communities, businesses, researchers, resource managers, international organizations and policymakers).

South Pacific and Southern Ocean

- This region consists of three main subregional fisheries: coastal South American pelagic fisheries, including the world's largest (Peruvian anchoveta); Antarctic krill, which has major development potential (biomass estimated at 500 million tons); and several demersal and pelagic species of the South-West Pacific, including blue grenadier, mackerel and squid.
- The highly productive Southern Ocean and the large upwelling zone off the coast of Peru and Chile have a naturally high carbon dioxide system which experiences periodic incursions of low pH water and may be particularly vulnerable to ocean acidification.
- Potentially deleterious effects from ocean acidification may impact key trophic links, such as shelled pteropods, which have already shown sensitivity to current carbon dioxide levels in the Southern Ocean.
- Sensitivity of molluscs to high carbon dioxide has been shown with effects of reduced growth, calcification and hatching and larvae recruitment success. Some aquaculture species or strains may be susceptible; however, others may have potential to acclimate or adapt.
- Aquaculture is prevalent in both the South-West and South-East Pacific. Key species are Atlantic and Coho salmon, rainbow trout, New Zealand and Chilean mussels, cupped oysters, calico scallops and *Gracilaria* seaweed. Ocean acidification could constrain future harvest of some species.
- Fisheries and aquaculture sectors are composed of a few large industrial fisheries, consolidated aquaculture and aqua-feeds industry, and a large number of small-scale fishers in Latin America.
- The limited number of key actors facilitates the possibility of dialogue and action to develop engagement strategies to target key fisheries, strengthen regional governance, foster collaboration among regional fisheries management

organizations, integrate actions with United Nations agendas, undertake selective breeding for increased resilience and explore risk analysis with the insurance industry.

North Atlantic and Arctic Ocean

- The North Atlantic has the highest proportion of the anthropogenic carbon distributed throughout the whole water column and the Arctic may experience corrosive surface waters in the near future.
- Ocean acidification may have pronounced interactive effects with strong background warming and expanding hypoxia.
- While ocean warming may increase general productivity in the North Atlantic, which may lead to an increase in fish catch, ocean acidification could constrain this increase in catch potential.
- Bivalves are likely to be impacted by ocean acidification, whereas effects on adult fishes may be small and indirect. The long-term resilience of cold water coral reefs as a nursery habitat for fishes is unknown.
- Fisheries and aquaculture in North America and coastal European countries are of major regional political, social and cultural importance and are a large fraction of the economies of Greenland, Iceland and the Faroe Islands.
- North America and Europe are substantial import markets for seafood and thus are dependent on ocean acidification impacts in other areas of the world.
- Ocean change due to warming and acidification has implications for production and trade patterns of natural wild (and invasive) fish stocks and aquaculture. Issues of industry adaptation, relocation and employment can be anticipated.

Central and South Atlantic

- This region includes subregions with special features, such as the eastern Atlantic high-productivity coastal upwelling areas (Guinea and Benguela currents) and the Caribbean Sea.
- Large upwelling areas are associated with high carbon dioxide waters. These conditions may become more extreme in the future as the ocean takes up more atmospheric carbon dioxide.
- Large rivers drain into this region, which are able to strongly alter the seawater pH near the shore beyond what may occur due to atmospheric carbon dioxide. The complex biogeochemical controls on seawater carbon dioxide chemistry make it difficult to detect trends of ocean acidification in coastal areas.
- Small-scale fisheries of small pelagics (for example, sardines, anchovies and herring) and some large-scale fisheries for demersal (hake), large pelagic (tuna) and small pelagic fish are found in all subregions of the Central and South Atlantic.
- The western Atlantic has major shellfish aquaculture activities, large-scale in the United States and artisanal in Brazil, which may be sensitive to ocean acidification.

- The sensitivity of open ocean finfish fisheries to ocean acidification via physiological or food-web effects is not yet known, but is of concern for coastal western and southern Africa.
- The Caribbean has important but degraded coral reef areas, which can be negatively impacted by ocean acidification. Alteration of the reefs will change the fish populations and regional productivity.
- Impacts that decrease harvests by small-scale fisheries have the potential to reduce food security and worsen food distribution inequalities that already exist in the region.

Mediterranean and Black Sea

- The coast is subject to strong human pressures with an estimated population of 132 million and intensive farming and industrial activities.
- Capture fisheries total 1.4 million tons landed each year, primarily small pelagic fish; marine and brackish aquaculture produces 180,000 tons of shellfish and more than 1 million tons of fish annually.
- In the Mediterranean Sea, fisheries and aquaculture provide 380,000 direct and 210,000 indirect jobs.
- Synergistic effects of warming and ocean acidification may increase the negative impacts on shellfish, especially early life stages, and other calcifiers such as red corals in this region. For aquaculture, recruitment and seed production are likely to be the main bottleneck for shellfish farming in the Mediterranean Sea.
- Impacts of ocean acidification on fish are not well studied, but may occur through changes in essential fish habitats or on other parts of the food web.
- Northern Mediterranean countries have more diversified fisheries-related economic activity, with higher production, consumption and export than southern Mediterranean countries, where production is largely intended for local consumption and export to the north.
- The socioeconomic impacts of ocean acidification will reflect the differences in economy and livelihood dependence on fish characterizing Mediterranean and Black Sea countries.

North and Central Pacific

- In many countries of this region, finfish and shellfish make significant contributions to government revenue, food security and employment. For example, the “coral triangle” is one of the world’s richest marine ecosystems, containing 75 per cent of reef-building species, and helps support over 120 million people with food, shore protection and income.
- Ocean acidification is already observable in this region; an open-ocean time-series station near Hawaii shows a decline in pH during the past 20 years.
- Upwelling regions along the west coast of the United States of America are particularly sensitive to increasing levels of carbon dioxide. Pacific deep waters

are becoming cumulatively corrosive. Washington state oyster aquaculture is the first example of ocean acidification affecting businesses.

- Carpet shell clams, cupped oysters, scallops, shrimp, sea cucumbers, coral reef fish and aquatic plants are all important marine harvest species in the region. Major capture fisheries include small (sardines, anchovy, mackerel and scads) and large (tuna) pelagic fish.
- The majority of laboratory studies indicate a negative impact of ocean acidification on bivalves; therefore, highly valuable aquaculture food enterprises producing clams and oysters in East and South Asia and scallops in China and Japan may be particularly sensitive.
- Despite the high economic value of pearl oysters in the tropical region, there is little work to date on the effects of ocean acidification on the production, quality and value of pearls from the black-lipped oyster. Data from a related species of pearl oyster suggest that shells are weakened by lower pH.

Indian Ocean and Red Sea

- More information is needed on carbon chemistry and fisheries in the Indian Ocean. Much of the wild fisheries catch is unreported or unidentified.
- Monsoon-induced upwelling along East Africa, the Arabian Sea and the Andaman Sea make these coastal waters seasonally high in primary productivity. This supports productive fisheries, but the upwelled high carbon dioxide water also make these areas particularly sensitive to ocean acidification.
- The commercial fish catch of the Indian Ocean region consists of tuna, sardines, mackerel, jacks, scads, pomfrets, kawakawa, octopus and squid. Primary shellfish harvests include clams, scallops, abalone, oysters and pearl oyster farms. Tropical coral reefs are found throughout the region. Ocean acidification impacts are likely to have negative effects on coral reefs, which may lead to altered reef fish habitat.
- The Indian Ocean region consists of mainland eastern and southern Africa, the Indian Ocean Island States, South Asia, and western and southern Australia. An estimated 800 million people live within 100 km of the coasts surrounding the Indian Ocean. Many members of coastal communities are undernourished, impoverished and highly dependent upon fishing for sustenance. Changes to harvest could therefore be a threat to food security.
- Most of the seafood harvests are small-scale artisanal fisheries. Negative effects of ocean acidification are anticipated for mollusc fisheries and mariculture. Aquaculture in sub-Saharan Africa and South Asia is increasing with large future potential. Shifts towards new production methods and cultured species may provide benefit to household livelihoods and small and medium enterprise development.

How does ocean acidification affect fisheries?

The oceans moderate the rate and severity of climate change by absorbing large amounts of carbon dioxide generated primarily from man-made activities. The uptake triggers large-scale changes in seawater chemistry, referred to as ocean

acidification due to the increase in seawater acidity (decrease of pH). In turn, these chemical changes alter physical and biological processes. Ocean acidification negatively affects many organisms that produce a calcium carbonate shell or skeleton, such as shellfish and corals. Other effects of high carbon dioxide include changes in growth rate, reproductive success or animal behaviour. The magnitude of the effect depends on the capacities of species to acclimate or adapt. Some organisms may be influenced by ocean acidification effects on components of the marine food web. For example, sea snails (shelled pteropods), which are food for salmon in the North Pacific, are threatened by acidification of high latitude waters. Effects are expected to be exacerbated when combined with other stressors, such as increased temperature, depleted oxygen and pollution. Ocean change may have major consequences for some keystone species, which may potentially lead to extinctions.

General information about ocean acidification and fisheries

- Knowledge of effects of ocean acidification on fish is currently inadequate.
- Effects of lowered pH on fish eggs and larval development have not been sufficiently studied.
- Coastal upwelling of deep ocean water to the surface can produce localized acidification.
- Impacts of ocean acidification on fisheries may include altered food webs.
- Small-scale artisanal fisheries employ 90 per cent of the world capture fishers and are important to food security and poverty alleviation (FAO).
- Nutrition from fish and seafood provides essential fatty acids, minerals and vitamins and is a primary source of protein for 1 billion people (FAO).
- Fisheries and aquaculture production, distribution and marketing employs 660 million to 800 million people, representing 10-12 per cent of the world population (FAO).

Filling the gaps in knowledge

Oceanographic and biological effects

- Effective monitoring of ocean acidification can establish clear cause and effect between changes in fisheries harvests and ocean acidification and provide early warning to fish farmers and fisheries managers.
- Ecosystem models need to be developed to project future acidification conditions in vulnerable and sensitive productive ecosystems.
- Reducing uncertainty by researching the effects of ocean acidification on seafood is necessary in order to evaluate the direct economic impacts on society. Priority areas of research include finfish, high-value crustaceans (shrimp, crab, lobster) and early life stages of seafood species.
- More research is needed on the combined effects of environmental factors on target organisms in order to gain a better understanding of the conditions that challenge natural communities.

- Little is known on how ocean acidification effects will alter marine food webs. Also, little is known about how ocean acidification will impact species in high-value coastal habitats, such as seagrass, mangroves and coral reefs.

Economic valuation and social impact

- Because the biological impacts are still poorly understood, assessing the impact on jobs and the economy is challenging. Case studies are to be undertaken on the economic and social impact of ocean acidification on fisheries for the species most vulnerable to ocean acidification, including cultured species and species important for marine leisure activities.
- Communities need to have a clear understanding of how ocean acidification may alter local economies and the possible scale of potential economic impacts based on assessment of exposure, sensitivity and adaptive capacity.
- Comprehensive risk assessments should be designed and implemented to prioritize adaptive responses.

Ocean acidification case study in aquaculture

In 2008, oyster hatcheries on the north-west coast of the United States were near collapse and a \$270 million industry employing over 3,200 people was under threat. Newborn oysters were dying off, with production levels reduced by 80 per cent and the cause of the losses was unknown. Research eliminated the bacterial pathogen *Vibrio* as the culprit. Instead, the oyster deaths were strongly linked to water quality, namely, ocean acidification. The hatcheries are open circulation systems supplied from the coastal ocean and the production failure was correlated to deep-ocean, corrosive waters brought to the surface. Working together, oyster businesses and scientists established actions to take. Production levels were restored to 70 per cent of normal by 2010. In response to the crisis, the state of Washington has budgeted \$3.3 million to regional efforts concerning ocean acidification in coordination with national agencies. Hatchery managers now monitor coastal waters to identify approaching upwelled water and take steps to protect their operations.

Participants and contributors

This brochure is the result of a joint effort by natural and social scientists from 19 countries, who met for two and a half days at the Oceanographic Museum of Monaco in November 2012.

Denis Allemand, CSM-Monaco; Edward Allison, UEA-UK; Andreas Andersson, SIO/UCSD-USA; Alexander Arkhipkin, United Kingdom; Bernard Avril, IMBER-Norway; Manuel Barange, PML-United Kingdom; John Baxter, SNH-UK; Johann Bell, SPC-New Caledonia; Richard Bellerby, NIVA-Norway; Gilles Boeuf, MNHN-France; Luke Brander, HKUST-Hong Kong; Tony Charles, St. Mary's U.-Canada; William Cheung, UBC-Canada; Mine Cinar, Loyola U. Chicago-USA; Joshua Cinner, JCU-Australia; Sarah Cooley, WHOI-USA; Ned Cyr, NOAA-USA; Cassandra DeYoung, FAO-Italy; Sam Dupont, U. Gothenburg-Sweden; Pierre Failler, CEMARE-United Kingdom; Laure Fournier, Total Foundation-France; Jean-Pierre Gattuso, CNRS-France; Frederic Gazeau, CNRS-France; Leigh Gurney, EC/JRC-Italy; Jason Hall-Spencer, Plymouth U.-United Kingdom; Lina Hansson, IAEA-Monaco; Gunnar Haraldsson, OECD-France; Nathalie Hilmi, CSM/IAEA-

Monaco; Courtney Hough, FEAP-Belgium; Christopher Kavanagh, IAEA-Monaco; Kieran Kelleher, Ireland; Thomas Lacoue-Labarthe, IAEA-Monaco; Dan Laffoley, IUCN-United Kingdom; Vicky Lam, UBC-Canada; Jean-Pierre Lozato-Giotart, Monaco; Fabio Massa, GFCM/FAO-Italy; Paula Moschella, CIESM; Paulo A.L.D. Nunes, CIESM; Laura Parker, UWS-Australia; Nicolas Pascal, CRIOBE-USR 3278 CNRS EPHE-Moorea; Gretta Pecl, UTAS-Australia; Hans O. Pörtner, AWI-Germany; Katrin Rehdanz, ifw/U. Kiel-Germany; Stephanie Reynaud, CSM-Monaco; Alain Safa, IDRAC-France; Melita Samoily, CORDIO-Kenya; Didier Sauzade, Plan Bleu-France; Juan Carlos Seijo, U. Marista de Merida-Mexico; Francois Simard, IUCN-Switzerland; Rashid Sumaila, UBC-Canada; Aurelie Thomassin, MEDE-France; Carol Turley, PML-UK; Michel Warnau, IAEA-Monaco; Wendy Watson-Wright, IOC/UNESCO-France; Patrizia Ziveri, UAB-Spain

Citation: Hilmi N, Allemand D, Betti M, Gattuso J-P, Kavanagh C, Lacoue-Labarthe T, Moschella P, Reynaud S, Warnau M (2013), Second International Workshop on the Economics of ocean acidification: bridging the gap between ocean acidification impacts and economic valuation, "Ocean acidification impacts on fisheries and aquaculture". Oceanographic Museum of Monaco, 11-13 November 2012.

Workshop websites:

<http://www.iaea.org/monaco/EconomicsOceanAcidification>

http://www.centrescientifique.mc/csmfr/informations/2012_11_OA.php

Acknowledgement: The workshop organizers extend their sincere thanks to all contributors to the production of this brochure, with special acknowledgement to Gretta Pecl, David Tarbath, Bruce Miller, Courtney Hough, Melita Samoily and George Waweru Maina for photographs and to Elsa Gärtner for the map.